


(Pdf free) A system of practical nosology; to which is prefixed, a synopsis of the systems of Sauvages, Linnus, Vogel, Sagar, Macbride, Cullen, Darwin, Crichton, Pinel, Parr, Swediaur and Young

A system of practical nosology; to which is prefixed, a synopsis of the systems of Sauvages, Linnus, Vogel, Sagar, Macbride, Cullen, Darwin, Crichton, Pinel, Parr, Swediaur and Young

David Hosack

*ebooks | Download PDF | *ePub | DOC | audiobook*



[Download](#)

[Read Online](#)

#12397374 in Books 2012-03-06Original language:EnglishPDF # 1 9.69 x .17 x 7.44l, .35 #File Name: 113067198480 pages | File size: 75.Mb

David Hosack : A system of practical nosology; to which is prefixed, a synopsis of the systems of Sauvages, Linnus, Vogel, Sagar, Macbride, Cullen, Darwin, Crichton, Pinel, Parr, Swediaur and Young before purchasing it in order to gage whether or not it would be worth my time, and all praised A system of practical nosology; to which is prefixed, a synopsis of the systems of Sauvages, Linnus, Vogel, Sagar, Macbride, Cullen, Darwin, Crichton, Pinel, Parr, Swediaur and Young:

This historic book may have numerous typos and missing text. Purchasers can download a free scanned copy of the original book (without typos) from the publisher. Not indexed. Not illustrated. 1821 Excerpt: ...by severe deep seated pain in the parts affected. The lymph of the vesicles, which is at first clear and colourless, becomes gradually milky and opaque, and ultimately concretes into scabs; but, in some cases, a copious discharge of it takes place, and tedious ulceration ensue. Not contagious. SPECIES OR VARIETIES. 1. Herpes phlyctodes. 2. zoster. 3. circinatus. 4. labialis. 5. ---praxialis, 6. iris. Genus IV. RUPIA. Characterized by an appearance of broad and flattened vesicles in different parts of the body, especially upon the extremities; they do not become confluent; they are slightly inflamed at the base, slow in their progress, and succeeded by an ill-conditioned discharge, which concretes into thin and superficial scabs; they are easily rubbed off, and presently regenerated. SPECIES OR VARIETIES. 1. Rupia simplex. 2., prominens. 3. ---escharotica. Genus V. MILIARIA. Miliaria, characterized by a scattered eruption of minute round vesicles, about the size of millet seeds, surrounded by a slight inflammation or rash, and appearing at an uncertain period of febrile disorders. The eruption is usually preceded by profuse perspiration, a sense of great heat, with a prickling and tingling in the skin. Most abundant upon the neck, breast, and back, sometimes in irregular patches, and sometimes more generally diffused. During the progress of the disease, aphthous vesicles and sloughs sometimes appear in the mouth and fauces. Genus VI. ECZEMA. The eczema is characterized by an eruption of small vesicles on various parts of the skin, usually set close or crowded together, with little or no inflammation round their bases, and unattended by fever. Not contagious. Unless extensively diffused, not accompanied with any derangement of the constitution. SPECIES OR V...